

System size and beam energy dependence of azimuthal anisotropy from PHENIX

Michael Issah (for the PHENIX Collaboration)

Department of Physics and Astronomy, Vanderbilt University, 1807 Station B, Nashville
TN 37235, USA

E-mail: michael.issah@vanderbilt.edu

Abstract.

We present azimuthal anisotropy measurements in Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 62.4$ and 200 GeV. Comparison between reaction plane and cumulant v_2 measurements in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV show that non-flow contributions, originating mainly from jets, influence the extracted v_2 for $p_T \gtrsim 3.5$ GeV/c. Number of constituent quark (NCQ) scaling of v_2 , when studied as a function of transverse kinetic energy KE_T , is seen to hold for Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ and 200 GeV and for Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV for $KE_T \lesssim 1$ GeV/c. Differential hexadecupole flow v_4 seems to exhibit scaling with integral v_2 for centrality $\leq 40\%$ as has been observed for differential v_2 .

1. Introduction

Experimental measurements at the Relativistic Heavy Ion Collider (RHIC) have indicated the creation of a deconfined phase of quarks and gluons [1]. The azimuthal anisotropy of particles emitted suggests that the high-energy density matter rapidly thermalizes and its evolution is driven by large pressure gradients [2, 3]. Azimuthal correlation measurements in Au+Au collisions at RHIC have been shown to consist of a mixture of jet and harmonic contributions [4]. The azimuthal anisotropy extracted from these correlations are thus the sum of contributions from collective flow and jet-like correlations. The latter constitute one source of non-flow correlations, especially at high p_T , and one would like to study how they vary with p_T and centrality. It has been reported that constituent quark scaling of v_2 as a function of KE_T or $m_T - m$ is observed in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and that differential v_2 scales with integral v_2 for both Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV [5]. These observations have implications on the hydrodynamic description of the medium as well as the pertinent degrees of freedom in the flowing matter [5]. PHENIX has measured the v_2 of charged pions, kaons and protons in Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV. It is interesting to study whether NCQ scaling holds for these two colliding systems and if the scaling of differential v_2 with integral v_2 also works for differential v_4 .

2. Data analysis

The results presented here use data obtained by the PHENIX collaboration for Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ and 200 GeV and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV. The azimuthal anisotropy relevant to this work were measured using two different methods: the reaction plane [6, 7] and the cumulant [8] method. The reaction plane method measures the correlations between the azimuthal angle of the particles detected in the central arms of the PHENIX detector and the azimuth of the reaction plane obtained with the two Beam-Beam Counters (BBC) located at $|\eta| \sim 3 - 3.9$ [9]. The cumulant method extracts v_2 from multiparticle correlations of particles detected in the PHENIX Central arms [10]. The large η separation between the BBC's and the central arms of PHENIX means that the extracted anisotropy is much less affected by possible non-flow contributions, especially those due to jets [9, 11]. The differential second-order cumulant v_2 is determined with respect to the reconstructed v_2 integrated over p_T . Monte Carlo simulations have been carried out to ensure that the azimuthal anisotropy is reconstructed faithfully with the PHENIX detector acceptance [8, 10]. The particles used to reconstruct the differential flow are excluded from those used in the integral flow reconstruction to avoid autocorrelations. The second order cumulant $v_2\{2\}$ is extracted from two-particle correlations and thus includes correlations from particles originating from jet fragmentation. Hence, the second order cumulant v_2 is influenced by non-flow correlations from jets as discussed in [8]. The results presented in the next section show how these jet correlations affect the azimuthal anisotropy in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

3. Results

3.1. Comparison between v_2 using different methods of measurement




Figure 1. Comparison between v_2 from cumulant and reaction plane (RP) methods for different centralities in Au+Au at $\sqrt{s_{NN}} = 200$ GeV. The closed and open circles, labelled $v_2\{RP\}$ and $v_2\{2\}$, are the cumulant and reaction plane results respectively.

Fig. 1 shows the differential v_2 in Au+Au at $\sqrt{s_{NN}} = 200$ GeV as a function of p_T from the reaction plane and the cumulant methods for several centralities. The measured

v_2 from the two methods agree well up to $p_T \approx 3.5$ GeV/c but differ for higher p_T , with the cumulant v_2 being higher than the reaction plane v_2 . This could be attributed to the influence of jets on the extracted v_2 which is more prominent in the cumulant method. One also notes that the same trend is observed at all centralities, even though the agreement between the two sets of results holds up to higher p_T for the 0-10% centrality bin.

3.2. Number of constituent quark scaling

Number of constituent quark scaling of v_2 in Au+Au at $\sqrt{s_{NN}} = 200$ GeV works well when studied as a function of transverse kinetic energy, as shown in the left panel of Fig. 2 [5]. This scaling scheme has been extended to the v_2 of positive and negative pions, kaons and protons measured in Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV. The middle panel of Fig. 2 shows that the constituent quark scaling of v_2 works well for $\pi^\pm$, $K^\pm$ and $(\bar{p})p$ in Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV. The same good scaling of v_2 is observed for the smaller Cu+Cu system at $\sqrt{s_{NN}} = 200$ GeV, as shown in the right panel of Fig. 2. These observations suggest that partonic degrees of freedom exist in the matter formed at lower beam energy and in the smaller colliding system.




Figure 2. v_2/n_q as a function of KE_T/n_q for charged pions, kaons and protons in minimum bias Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV (left panel), in Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV for centrality 10 - 40% (middle panel) and Cu+Cu at $\sqrt{s_{NN}} = 200$ GeV for centrality 0 - 50% (right panel).

3.3. v_4 of unidentified particles

PHENIX has recorded high statistics Au+Au collisions during the 7th year of data-taking, enabling the detailed study of higher harmonics of azimuthal anisotropy as a function of p_T and centrality. It has been reported that v_2 scales with the integral flow, which has been shown to be proportional to the eccentricity [5]. The use of the integral flow scaling of differential v_2 has been used to study the thermalization of the matter at RHIC [5]. Since, in general, $v_n \sim v_2^{n/2}$, which yields $v_4 \sim v_2^2$ for $n = 4$, differential v_4 would be expected to scale with the square of the differential v_2 and the scaling of differential v_4 with integral v_2 could be investigated. The left panel of Fig. 3 shows the p_T dependence of v_4 of unidentified particles for three centrality bins in the range 10-40 % and the right panel shows v_4 , scaled

by the square of the p_T -integrated v_2 for the corresponding centrality (labelled ϵ' in the figure), as a function of p_T . The latter plot shows that v_4 scales with the square of integral v_2 for centrality $\leq 40\%$ and $p_T \lesssim 2.0$ GeV/c. By plotting $(v_2/\epsilon')^2$ on the same plot (open circles) one observes a good agreement between differential v_4 and v_2^2 , when they are both scaled by the square of integral v_2 .




Figure 3. Left panel: Differential v_4 of unidentified hadrons as a function of p_T (left panel) for different centralities. Right panel: v_4/ϵ'^2 as a function of p_T , where ϵ' represents v_2 integrated over p_T . For the sake of comparison, the open circles represent differential v_2 squared scaled by the square of the integral v_2 .

4. Summary

In summary, we have presented results of azimuthal anisotropy from different methods at RHIC. They show that non-flow correlations from jets may influence the measured anisotropy for $p_T \gtrsim 3.5$ GeV/c. Constituent quark scaling is observed to hold in Au+Au at $\sqrt{s_{NN}} = 62.4$ and 200 GeV and Cu+Cu at $\sqrt{s_{NN}} = 200$ GeV. This suggests that partonic degrees of freedom are manifest at the lower RHIC beam energy and in the smaller Cu+Cu colliding system. The differential hexadecupole flow v_4 in Au+Au at $\sqrt{s_{NN}} = 200$ GeV is observed to scale with the integrated v_2 for centrality $< 40\%$ and $p_T \lesssim 2.0$ GeV/c.

References

- [1] K. Adcox *et al.* [PHENIX Collaboration], Nucl. Phys. A **757**, 184 (2005)
- [2] M. Gyulassy and L. McLerran, Nucl. Phys. A **750**, 30 (2005) [arXiv:nucl-th/0405013].
- [3] U. W. Heinz, J. Phys. Conf. Ser. **50**, 230 (2006) [arXiv:nucl-th/0504011].
- [4] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **90**, 082302 (2003) [arXiv:nucl-ex/0210033].
- [5] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 162301 (2007) [arXiv:nucl-ex/0608033].
- [6] J. Y. Ollitrault, Phys. Rev. D **48**, 1132 (1993) [arXiv:hep-ph/9303247].
- [7] A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C **58**, 1671 (1998) [arXiv:nucl-ex/9805001].
- [8] N. Borghini, P. M. Dinh and J. Y. Ollitrault, Phys. Rev. C **64**, 054901 (2001) [arXiv:nucl-th/0105040].
- [9] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 182301 (2003) [arXiv:nucl-ex/0305013].
- [10] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **94**, 232302 (2005) [arXiv:nucl-ex/0411040].
- [11] A. Adare *et al.* [PHENIX Collaboration], arXiv:0801.4545 [nucl-ex].